

Growth, Yield and Economics of sweet corn (*Zea mays* L) as influence by different Integrated Nutrient Management practices

S. KRISHNAPRABU

Department of Agronomy, Faculty of Agriculture, Annamalai University.

Received 14 June 2015 • Revised 23 August 2015 • Accepted 24 September 2015

Abstract: Field experiment was conducted at Department of Agronomy, experimental farm at Annamalai University during 2013. The main objective of the study was to find out the effect of organic & inorganic fertilizer on seed yield of sweet corn (*zea mays* L.). The experiment was studied with split plot design having two levels of biofertilizer, FYM and phosphorus as main plot treatments along with five levels of nitrogen as sub plot treatment. Application of organic matter had significantly increased height & all crop growth parameters and 5.75% more grain yield with application of FYM@10 ha⁻¹ Seed inoculation with *Pseudomonas* gave significant increase in growth and yield parameters and grain yield increased to the tune of 8.24 per cent.

Application of phosphorus significantly increased plant height at all crop growth stages and higher grain yield recorded by 6.74 per cent than Control. Seed yield of sweet corn as well as growth and yield attributes were significantly increased due to varying levels of nitrogen. The higher grain yield (1633 kg ha⁻¹) and straw yield (5783 kg ha⁻¹) was recorded with 120 and 160 kg N ha⁻¹ respectively. The net realization of Rs. 30525 & 29255 ha⁻¹ was recorded with 101 FYM ha⁻¹ Seed inoculation gave 31485 Rs. ha⁻¹ and application of P205 at 0 and 50 kg. P205 ha⁻¹ gave 6.70% and 4.43% higher grain and straw yield, respectively. The significant higher straw yield were recorded with varying levels of Nitrogen, but highest net return was obtained with 120 kg nitrogen per hectare, The straw yield were increased significantly with increasing levels of nitrogen from 0 to 160 kg per hectare.

Keywords: Economic, FYM, INM, organic fertilizer, sweet corn, *Zea mays*.

INTRODUCTION

Maize (*Zea mays* L.) popularity known as corn is one of the most important cereal of the world, ranking third amongst the food crops, next to rice and wheat both in respect of area and production. India occupied an area of 10.58 lakh hectares with the production of 14.32 lakh tones during the year 1993 correspondingly the Gujarat State had an area of 3.68 lakh ha with the production of 5.29 lakh tonnes. In Gujarat, maize is one of the important traditionally grown crop of tribal areas, comprising the districts of Panchmahals, Sabarkantha, Banaskantha and part of Baroda & Kheda districts, Now recently this crop may be introduced in South Gujarat districts like Surat & Tapi. Among these districts Panchmahals is a leading district which accounts for area of 2.62 lakh hectares and production of 2.15 lakh tonnes.

Among various types of maize, sweet corn is very popular for the use of its green cobs in the United States of America. It differs from the field corn due to its higher sweetness, as it has high amount of sugar & alcoholic material. Besides, its consumption as vegetable purpose, it is also utilized for extracting sucrose as an industrial purpose. The role of O.M. for increasing crop production has been universally established, as it plays significant role in improving physical and chemical properties of the soil. Application of 12-15 tonnes of FYM helps in increasing the yield of maize crop to the tune of 1.5 to 5.6 qa

/ ha. Sweet corn is one of the heavy consumers of plant nutrients. It remains about 72 kg N, 25 kg P₂O₅ and

220 kg K₂O / ha. Nitrogen is the key element in crop growth and is the most limiting nutrient in Indian soils. The importance of nitrogen for increasing the yield has been widely accepted.

Maize is one of the crop that responses well to phosphoric fertilizer in almost all the soil types. It plays vital role in plant nutrition. The deficiency of phosphorus is soil severely limits root and shoot growth and thereby affecting the yield. The availability of phosphorous are also low as compared to that of N & K. Under such situation, the phosphate solubilizing micro organism plays significant role in making the phosphorous available to plants by secretion of organic acids and enzyme phosphatase which solubilizes the insoluble phosphate and thereby it helps in increasing the crop production.

MATERIAL AND METHODS

Field experiment was conducted at Department of Agronomy, experimental farm at Annamalai University during 2013. The experiment was laid out on loam soil, clay well drained & fairly moisture retentive but low as compared to black soil. The experiment was laid out in forty treatments comprising all possible combinations of two levels of O.M. (FYM), two levels of *Pseudomonas*, two levels of phosphoric along with five levels of nitrogen. The study was carried out with split plot design (SPD). Combination of FYM x inoculations x phosphorous were taken as main plot treatment while levels of nitrogen were taken as subplot treatments with three replication having 5.4 m x 3.6 m gross plot size, 60 cm x 20 cm spacing and dibbling method of sowing. Application of well decomposed FYM as basal at 101 ha⁻¹ as per treatment 20 % of total nitrogen of respective levels of N compiled with full dose of phosphorous in form of SSP in a previously open furrow at the depth of 8-10 cm. Remaining 80% of nitrogen was applied in two installments UBC 50% of the total quantity at knee height stage and remaining 30% N of total amount at milking stage, The observations were recorded from five randomly selected plants from net plot (plant height, no. of barren plants) on growth & yield attributing character and also economics of (length of cob, number of cobs per plant, kernels row per cob, no. of kernels per cob, grain & straw yield) of sweet corn.

RESULT & DISCUSSION

Growth and yield: With a view to study the effects of organic matter, inoculation of phosphate solubilizing microorganism along with levels of phosphorous and nitrogen on the growth and grain yield of sweet corn (*Zea mays* L.).The findings on the yields of growth and yield attributed characters and economics as influenced by different treatments are showed in Table 1 and 2. Effect of seed inoculation with *Pseudomonas* was found significant in respect to plant height at all growth stages. Seed treated with *Pseudomonas* gave significant taller plants as compared to un inoculated seed. Measured at 21 days interval, i.e. 21,42.63 and 84 DAS at all growth (16.44, 46.72, 137.33, 143.09) and grain (1340 kg ha⁻¹) stages this might be due to the ability of Phosphobacteria to bring soluble / insoluble inorganic and organic phosphates into soluble forms by secretion of organic acids. Similar results were also noted by Kataraki et al. (2004). Application of FYM on sweet corn found non-significant effect of FYM on plant height measured periodically at 21, 63 & 84 DAS. However, the application of FYM gave numerically higher values of plant height at each period of crop growth stage and higher grain yield 1325 kg ha⁻¹, but straw yield had non significant effect. This could be attributed to the lower mineralization of organic nitrogen. Such observation was also made by Sahoo and Mahapatra (2004).

Table 1. Effect of integrated nutrient management of growth of sweet corn as influenced by levels of inoculation, FYM, phosphorus & nitrogen levels

Treatment	Plant height (cm)			
	21 DAS	42 DAS	63 DAS	84 DAS
Inoculation				
C ₀ uninoculated	15.29	40.91	129.38	142.18
C ₁ Inoculated	16.44	46.72	137.33	143.63
S.E.M +	0.07	0.26	0.34	0.41
CD at 5%	0.21	0.80	1.02	1.24
FYM t ha ⁻¹				

F ₀ 0	15.79	42.57	133.21	142.49
F ₁ 10	15.99	45.03	133.50	143.32
S.E.M ±	0.07	0.26	0.37	0.41
CD at 5%	NS	0.80	NS	NS
Phosphorus kg ha ⁻¹				
P ₀ 0	15.58	43.02	131.04	141.04
P ₁ 50	16.25	44.58	135.68	144.78
S.E.M ±	0.07	0.26	0.34	0.41
CD at 5%	0.21	0.80	1.02	1.29
C.V. %	3.38	4.63	1.95	2.22
Nitrogen kg ha ⁻¹				
N ₀ 0	14.69	37.86	122.75	135.98
N ₁ 40	15.30	40.04	129.96	139.21
N ₂ 80	16.12	44.50	135.63	142.80
N ₃ 120	16.24	47.00	188.33	146.63
N ₄ 160	16.96	49.58	140.04	149.94
S.E.M +	0.10	0.48	0.65	0.53
CD at 5%	0.28	1.38	1.86	1.51

Application of phosphorus @ 50 kg P₂O₅ ha⁻¹ was found significant on the plant height measured at all growth stages, grain yield (1331), straw yield 4777 kg ha⁻¹ (*i.e.* 21,42.03 & 84 DAS) significant increase plant height (16.25, 44.58, 135.68,144.75) due to the phosphorus as a key element influences different physiological process such as cell division and elongation (Pandey *et al.*, 2000).

Significant linear increase in plant height, grain & straw yield was observed with each successive increase in N levels from 0 to 160 kg ha⁻¹. Significantly increased the grain yield (1331 kg ha⁻¹) & straw yield (5783 kg ha⁻¹) with 160 kg ha⁻¹ and also showed the maximum plant height (16.93, 49.58, 140.04, and 149.92) at all the crop growth stages. The higher availability of nitrogen might have increased its uptake as a results of which increased cell size and enhanced cell division, seems to have played an important role in increasing the plant height and yield, this findings confirms to those reported by Sharma and Gupta (1998). Interaction effect between inoculation and phosphorus, FYM with phosphorus and nitrogen levels all the interaction effect were significantly gave higher growth of and grain & strover yield of sweet corn.

Table 2. Grain, straw yield & economics of sweet corn as influenced by inoculation, FYM, phosphorus and nitrogen levels

Treatment	Grain yield real yield (kg ha ⁻¹)	Straw yield (kg ha ⁻¹)	Gross realization (Rs./ha)	Total cost of cultivation (Rs./ha)	Net realization (Rs./ha)	BCR
Inoculation						
C ₀	1241	4481	32745	3900	28845	7.39
C ₁	1340	4838	35435	3950	31485	7.97
S.E.M ±	20.14	62.54				
CD at 5%	61.08	189.61				
FYM (t ha ⁻¹)						
P _Q	1256	4575	33155	3900	29255	7.50
F _i	1325	4752	35025	4504	30525	6.78
S.E.M +	20.14	62.51				
CD at 5%	61.08	NS				
Phosphorus						
F ₀	1249	4550	32995	3900	29095	7.48
F ₁	1331	4777	35185	4540	30645	6.75
S.E.M +	20.14	62.51				

CD at 5%	61.08	189.61				
C.V. %	12.10	10.38				
Nitrogen						
N ₀	1015	3447	26703	3900	22814	5.86
N ₁	1128	3935	29774	4272	25502	5.96
N ₂	1341	4671	35371	4543	30828	6.78
N ₃	1633	5481	43017	4815	38202	7.93
N ₄	1334	5783	35588	5087	30501	5.94
S.E.M ±	23.82	78.69				
CD at 5%	67.38	222.25				
Interaction						
CXP						
c.v.%	9.01	8.29				

ECONOMICS

The data on gross and net realization for different treatments of FYM, inoculation, phosphorus and nitrogen presented in Table 2 revealed that the higher net returns of Rs. 30525 / ha were received with treatment F, (FYM 50kg/ha) of the seed treatment with *Pseudomonas* sp. gave a higher net returns & Rs. 31485/ha as compared to uninoculated control. Application as 50 kg P₂O₅ / ha gave a higher net return of Rs. 30645 / ha as compare with no application of phosphorus. Among the nitrogen levels, the maximum net returns of Rs. 38202/ha were realized with the application of 120 kg ha⁻¹. The results confirm the findings of Adhikari *et al.* (2005).

CONCLUSION

On the basis of study the results obtained from the investigation the conclusion can be draw for getting maximum seed and thereby net monetary realization the sweet corn should be fertilized with 101 FYM + 120 kg N ha⁻¹ + 50 kg P₂O₅ besides, seed inoculation with *Pseudomonas* sp.

REFERENCES

- [1]. Adhikari, S., Chakraborty, T. and Bagchi, D.K. 2005. Bio- economic evaluation of maize (*Zea mays* L.) and groundnut (*Arachis hypogaea*) intercropping in drought prone areas of Chotonagpur Pleateau region of Jharkhand. *Indian Journal of Agronomy*. 50 (2): 113-115.
- [2]. Jadhav V, T and D.K. Shelke 2012. Effect of Planting methods and fertilizer levels on Growth, yield and Economics of maize. hybrids *J. of Agri, Res and Technology*. 37(1): 11-14.
- [3]. Kataraki, N.G., Desai, B.K. and Pujari, B.T. 2004. Integrated nutrient management in irrigated maize. *Karnataka Journal of Agriculture Sciences*. 17(1) : 1 -4.
- [4]. Kumar P., A.S. Halepyati, B.T. Pujari and B.K. Desai 2007. Effect of INM on Productivity, nutrient uptake and economics of maize under reinfed condition. *Karnataka J. of Agri Sci.*, 20 (3): 462-465.
- [5]. Massey J.X and B.L. Gaur 2006. Effect of Plant population and fertilizer levels in growth and NPK uptake by sweet corn *Ann. Of Agri. Res*. 27(4): 365-368.
- [6]. Pandey A.K., V.P. mani, Ved Prakash, R.D. Singh and H.S. Gupta 2002. Effect of varaieties and Plant densities on yield, yield attributes and economics of baby corn.
- [7]. Pandey, A.K., Ved, Prakash, Mani, V.P. and Singh, V.P 2000: Effect of rate of nitrogen and time of application on yield and economics of baby corn (*Zea mays* L.) *Indian Journal of Agronomy*. 45 (2): 338-343.
- [8]. Sahoo, S.C. and Mahapatra, P.K. 2004. Response of sweet corn (*Zea mays* L.) to nitrogen levels and plant population. *Indian Journal of Agricultural Sciences*. 74 (6): 337-338.
- [9]. Sharma, M.P. and Gupta, G.P. 1998. Effect of organic materials on grain yield and soil properties in maize - wheat cropping system. *Indian Journal of Agricultural Sciences*. 68 (11) : 715-717.
- [10]. Singh. U., A.A. Saad, T. Ram and L. chaud 2012. Productivity, economics amnd Nitrogen use efficiency of Sweet corn as influenced by planting geometry and N fertilization. *Indian J. of Agron* 57(1): 43-48.

- [11]. Thakur G. D., P. N. Karanjikar and A.B. Kasbe 2010. Effect of Fertilizer levels on yield and yield contributing characters of Sweet corn. Asian J. Of Soil Sci. 4(2): 280-282.
- [12]. Verma, N, K. 2011. INM in winter maizer sown at differnet rates, J. of plnat Breeding and Crop Sci., 3(8): 161-167.