

# Preparation and diagnosis of new dyes for pyrimidine-2, 4, 6(1H, 3H, 5H)-trione

Ali Ghazi Kadhim<sup>1</sup> - Ali Noory Fajer<sup>2</sup>

<sup>1</sup>AL-Qadisiyah University - Chemistry Dep. College of Education-

<sup>2</sup>Department of Biology, College of Education, University of Al-Qadisiyah, AlDwaniyah, Qadisiyyah, Iraq

E.MAIL: <sup>1</sup>ali.kazim@qu.edu.iq, <sup>2</sup>ali.fajer@qu.edu.iq

Received 18 November 2018 • Revised 18 December 2018 • Accepted 19 January 2019

**ABSTRACT:** The research includes the preparation of new azo-barbiturate dyes derived from Schiff bases prepared by reaction of barbituric acid & phenylenediamine and repellency bases by means of double-reaction of repellent bases with different phenolic compounds. These compounds were diagnosed using infrared techniques, ultraviolet-visible radiation, NMR proton and thin layer chromatography. All barbiturate dyes had absorption between (273 – 310) nm and showed hypsochromic shift from the corresponding azo dyes. The crystal structures of barbiturate dyes indicated that the barbiturate ring is sterically hindered by phenol rings.

**Keywords:** Barbituric acid, thymol, quinol, phenylenediamine.

## INTRODUCTION

Barbiturates derived from an acidic basis are barbituric acid (**1**) and are characterized as medium to low acidity compounds with acidic function (7-8) (**2**). These compounds are not insoluble in water (**3**) and have been prepared to overcome this characteristic by reacting with sodium ions, which is prepared in the form of salts and becomes soluble in water (**4**). Several barbiturates were derived by substituting the roots in their basic construction, barbituric acid, which scientists agreed to number their atoms (**5**). These derivations include the radical substitution of ethyls where hydrogen atoms on carbon 5 resulted in the discovery of the barbital compound (**6**). The replacement of a vinyl root by the location of one of the ethyl radicals on carbon 5 (in the barbital compound) led to the discovery of phenobarbital, which is characterized as antimicrobial (**7**) and muscle dysfunction (**8**). Replace the sulfur element with the oxygen element on carbon 2 and replace the side chain with the place of one of the ethyl radicals on carbon 5 (in the barbital compound), resulting in the thiopantone compound (**9**). The radical replacement of the instance of the two hydrogen atoms associated with the azotone 1, 3 and the replacement of cyclohexanyl and the place of one of the ethyl radicals on carbon number 5 resulted in the acquisition of sodium hexobarbital (**10**). The latter two have the fastest impact, i.e., rapid anesthesia and short impact duration (**11**).

Schiff base a group called azomethine (C = N-) (**12 - 14**). Also, Schiff bases derived from aldehyde condensation with primary amines are called al-deamins. In the so-called Schiff bases derived from the condensation of ketones with the primary amines are called al-katamines (**15**). The stability of Schiff bases depends on the type of amine and the type of aldehyde or ketone used. The bases of the Schiff prepared from the aromatic ketone and the aromatic amine are the most stable between the Schiff bases. This is due to increased stability by resonance (**16**). Azomethine dyes are of great importance and wide use in industries Paper, colored reagents for food and cosmetics (**17-21**). Azomethine compounds are also bioactive, used as antimicrobial agents, cancer (**8**) and others (**22-25**).

## METHODS OF WORK

All compounds were purified by re-crystallization followed by thin layer chromatography and modern spectral techniques. As for melting points, it was measured using Staurat MP / MP3. Infrared spectra were recorded using a spectrometer (Shimadzu FT-IR 8400 S) with a potassium bromide tablet. Absorption spectra were measured in ethanol using spectroscopy (Shimadzu UV-Vis. 1600) at Qadisiyah University - Faculty of Education. The proton NMR spectrum was taken using a BRUKER (300 MHz)

spectrometer in Jordan - Al-Bayt University. Chromatography was measured using Silica Gel TLC plates 60 F250. (26-28)

### METHODS OF PREPARATION

The general method for preparing the base of azomethi -

In R.B.flask (100 ml), put 0.02 ml of PYRIMIDINE derivative and add 25 mL of pure ethanol and then (1 ml) gla. acetic acid, and gradually add (0.06 mol) aromatic amine with (20 ml) pure ethanol and then the mixture reflux for (2hr.) The reaction was followed by chromatography of the thin layer. After heating, the reaction mixture was cooled, the precipitate was then filtered and dried and then recrystallized using ethanol.

### The general way to prepare azo pigments


Preparation of diazotium salt sol. In a glass baker, prepare a solution consisting of 10 mL cooled water and 10 mL concentrated hydrochloric acid and with continuous stirring (0.02 mL) of the prepared shiff base in the first part (A) with cooling in the ice bath (0-5) M, then add to it a coolant solution consisting of (0.06 mol) sodium nitrite + (5 ml) water.

-In a second glass baker and within the ice bath, a solution consisting of dissolving (0.06 mol) phenolic compound with (15 ml) sodium hydroxide solution (10%).

After continuous stirring in the ice bath, add the diazonium salt solution and drops to the coupling solution. After the addition of the mixture, stir the reaction mixture for 15 minutes in the ice bath. The deposited precipitate is washed with water and then recrystallized using ethanol.


### RESULTS AND DISCUSSION (29-34)

This research included two steps: The first step was to prepare a new shiff base (A) by reacting the condensation between barbituric acid and the 1,4-phenylene diamine compound, scheme number(1).



scheme (1)

The second step involved the preparation of new azo dyes through the interaction of shiff base prepared in the first step using diazotization reaction with several phenolic compounds (**scheme 2**). Eight new azotecin derivatives were obtained (COMP.: A-H). Table (1) represents all physical properties of prepared derivatives. Table (2) represents all the spectral properties of the prepared compounds (18). Spectra FT-IR shows in Figures ((A - H)).



PhOH = PHENOL, 1-Naphthol , 2-Naphthol , Quinol ,Resorcinol , 2-aminophenol , thymol ) .

**scheme (2)**

**Table 1:** Physical properties of prepared derivatives (L:Light , D:Dark)

Com.	Phenol	M.Str.	M.wt.(g.mol <sup>-1</sup> )	Yield (g, %)	M.P.(°C)	Color
A	-----	C <sub>22</sub> H <sub>22</sub> N <sub>8</sub>	398.39	(0.3 , 80)	190-192	D.Orang
B	phenol	C <sub>40</sub> H <sub>31</sub> N <sub>11</sub> O <sub>3</sub>	713.74	(0.4 , 60)	205-206	yellow
C	1-Naphthol	C <sub>52</sub> H <sub>37</sub> N <sub>11</sub> O <sub>3</sub>	863.92	(0.5 , 70)	257-258	L. yello.
D	2-Naphthol	C <sub>52</sub> H <sub>37</sub> N <sub>11</sub> O <sub>3</sub>	863.92	(0.6 , 68)	271-273	Orange
E	Quinol	C <sub>40</sub> H <sub>31</sub> N <sub>11</sub> O <sub>6</sub>	761.74	(0.4 , 58)	232-234	Brown
F	Resorcinol	C <sub>40</sub> H <sub>31</sub> N <sub>11</sub> O <sub>6</sub>	761.74	(0.4 , 65)	230-232	D.yello.
G	2-Amino phenol	C <sub>40</sub> H <sub>34</sub> N <sub>14</sub> O <sub>3</sub>	758.78	(0.3 , 55)	240-241	L.orang.
H	Thymol	C <sub>52</sub> H <sub>55</sub> N <sub>11</sub> O <sub>3</sub>	882.06	(0.2 , 50)	265-267	D.brow.

Table 2: Spectral properties of prepared derivatives




Figure. A




Figure.B

Com.	Phenol	$\lambda_{\max}$ EtOH	FT-IR( $\text{cm}^{-1}$ ) KBr disk				
			- NH	-C=N	N=N	C-O	-OH ring
A	-----	-----	3373	1635	-----	1284	-----
B	phenol	273	3345	1631	1523	1290	1474
C	1-Naphthol	288	3362	1614	1522	1294	1478
D	2-Naphthol	290	3359	1618	1518	1293	1476
E	Quinol	275	3387	1616	1522	1291	1475
F	Resorcinol	278	3395	1617	1524	1294	1476
G	2-Amino phenol	295	3372	1620	1526	1296	1480
H	Thymol	310	3422	1623	1530	1298	1481




Fig c





Fig D



E Fig

Fig F

### Spectrometry (<sup>1</sup>H-NMR) of prepared derivatives

<sup>1</sup>H NMR (300MHz, CDCl<sub>3</sub>, δ / ppm):

**COMP.((A))** : 2.96 (d, 2 H, *J* = 14.8 Hz), 3.59 (d, 2 H, *J* = 14.8 Hz), 6.98 (m, 7 H), 7.06 (s, 5 H), 7.17 (m, 7 H).

**COMP.((B))** : 3.31 (s, 2 H), 6.96-7.03 (m, 8 H), 7.32 (m, 8 H), 7.50 (q, 7 H, *J* = 6.4 Hz), 7.60 (dd, 7 H, *J* = 7.0, 1.7 Hz).

**COMP.((C))** : 3.29 (s, 2 H), 6.96 (dd, 3 H, *J* = 8.6, 1.9 Hz), 7.22 (m, 7 H), 7.50-7.77 (m, 18 H), 7.81-7.88 (m, 3 H), 8.18-8.27 (m, 3 H).

**COMP.((D))** : 3.29 (s, 2 H), 6.98 (d, 2 H, *J* = 8.8 Hz), 7.04 (d, 1 H, *J* = 8.6 Hz), 7.22 (m, 7 H), 7.50-7.70 (m, 15 H), 7.76 (dd, 1 H, *J* = 8.2, 1.6 Hz), 7.86-8.00 (m, 8 H), 8.11 (dt, 1 H, *J* = 7.8, 2.1 Hz).

**COMP.((E))** : 3.23 (s, 2 H), 6.45-6.48 (m, 4 H), 6.83-6.87 (m, 4 H), 7.23 (m, 3 H), 7.31 (s, 11 H), 7.44-7.50 (m, 4 H), 7.67 (d, 3 H, *J* = 8.1 Hz).

**COMP.((F))** : 3.08 (d, 1 H, *J* = 15.3 Hz), 3.38 (d, 1 H, *J* = 15.3 Hz), 6.45-6.48 (m, 4 H), 6.83-6.87 (m, 4 H), 7.23 (m, 3 H), 7.31 (s, 11 H), 7.44-7.50 (m, 4 H), 7.67 (d, 3 H, *J* = 8.1 Hz).

**COMP.((G))** : δ 3.21 (s, 3 H), 6.48 (d, 1 H, *J* = 1.8 Hz), 6.55 (d, 3 H, *J* = 1.1 Hz), 6.68-6.71 (m, 4 H), 7.15 (m, 6 H), 7.21-7.31 (m, 12 H), 7.54 (d, 4 H, *J* = 8.2 Hz).

**COMP.((H))** : 1.32 (d, 20 H, *J* = 6.8 Hz), 2.08 (s, 3 H), 2.13 (s, 6 H), 3.27 (s, 2 H), 3.38 (h, 3 H, *J* = 7.1 Hz), 6.68 (s, 2 H), 6.80 (s, 1 H), 7.20-7.29 (m, 7 H), 7.46-7.51 (m, 9 H), 7.63 (s, 1 H).

The NMR spectrum of the azoketimines indicates that there are several overlapping electronic environments of the prepared compounds due to the diversity of the active groups found in these compounds in that they are driving or drawing aggregates (such as azomethane, hydroxyl, amine and alkyl groups), in addition to the abundance and diversity of hydrogen atoms (6.60 ppm) refers to the adjacent protons of the base aromatic ring, whereas the signal (6.80 ppm) belongs to the opposite protons of the aromatic compound. The opposite protons of the aromatic amine compound, close to the azomethin group, show a clear signal with double bursts of 7.81 ppm. For the latter, they are due to the opposite protons of the base aromatic ring, especially near the fivering.

For the phenol compound, one of the most significant signs of the characteristic resonance characteristic is the signal (6.99 ppm) of the corresponding phenol ring protons, which at the same time are protons adjacent to the other phenol loop protons, causing double and double fissures. As for the naphthol dye, the most important signals are measured between the extended values (ppm 7.09 - 8.03), which are related to the protons of the alpha-naphthol ring, which are close to the hydroxyl driving group and the torso-tzo group, Hydrogen atoms. As for azumethane and pyrethylene, the presence of the high-density electron groups creates electronic environments that obscure or reduce the appearance of signals and thus reduce their measured values or observations. This is observed for these compounds, the most important of which are the measured values 7.17 - 6.91 ppm), which is due to the proton ring between the hydroxyl driving groups. The characteristic and phenolic value of the phenolic compound is a signal between 7.18 - 6.80 ppm, which comes from protons confined between the iso group and the amino and

hydroxyl groups. The most recent of these thymol has several distinctive signs, the most important of which is the signal (3.23 ppm) of proton, the isopropyl group, which is affected by the high electronic environment of the two kinetic groups, as well as the signal (1.96 ppm) of the three methyl groups of the thymol. (Ppm 1.18). The signal (6.99 ppm) belongs to the thymol group, which is bound to the methyl and hydroxyl groups.

## CONCLUSIONS

The main objective of this research is to prepare new azotecins containing several phenolic derivatives using one of the important organic reactions, which is the reaction of azuna. The classical and modern techniques were used in the diagnosis of prepared compounds such as melting grades, chromatography, Ultraviolet-visual radiation, proton spectroscopy and NMR spectroscopy. The percentages of derived derivatives approximately good. We hope that our research will be done in the form of new azotecin dyes used in some industrial and applied fields such as dyes, various medicines, cosmetics, colored reagents and analytical chemistry guides .

## REFERENCES

- [1]. B. Dickey & A. R. Gray. "Barbituric acid". Org. Synth.,(1943), Coll. Vol., 2, p. 60 .
- [2]. .Kauffman, George B. "Adolf von Baeyer and the naming of barbituric acid," Journal of Chemical Education, (1980), 57 : 222-223.
- [3]. Boyd E M, Pearl M. "Can nalorphine hydrochloride prevent respiratory depression and death from overdose of barbiturates" , Canadian Medical Association Journal, (1955), 73(1):35-8.
- [4]. Koppanyi T, Fazekas J F. "Acute Barbiturate Poisoning Analysis and Evaluation of Current Therapy[J]. American Journal of the Medical Sciences", (1950), 220(5):559-576
- [5]. Shulman A, Shaw F H, Cass N M, et al. "A New Treatment of Barbiturate Intoxication"]. British Medical Journal, (1955), 1(4924):1238-44.
- [6]. Bateman C H. "BARBITURATE POISONING" J. Lancet, (1963), 282(7303):357.
- [7]. Ali.G. kadhim , " synthesis of new azo dyes via suzuki coupling ", (2014) .college of science 12 (20), 66-72
- [8]. "Barbiturates". Archived from the original on 7 November 2007. Retrieved 2007-10-31, "Dorlands Medical Dictionary:hypnotic" . Mercksource.com. Archived from the original on ,(2008),12-11.
- [9]. Brunton, Laurence L.; Lazo, John S.; Lazo Parker, Keith L.. "17: Hypnotics and Sedatives". Goodman & Gilman's The Pharmacological Basis of Therapeutics", (2006) , (11th ed.).
- [10]. Mendels, J. "Criteria for selection of appropriate benzodiazepine hypnotic therapy". (September 1991). J Clin Psychiatry. 52. Suppl: 42-6
- [11]. Glass, J.; Lanctôt, K. L.; Herrmann, N.; Sproule, B. A.; Busto, U. E. "Sedative hypnotics in older people with insomnia: meta-analysis of risks and benefits".(November 2005). BMJ. 331 (7526).
- [12]. Arulmurugan S, Kavitha PH, Venkatraman RP . " Biological activities of Schiff base and its complexes: a review ". Rasayan J Chem . ,(2010) , 3(3):385-410.
- [13]. Przybylski P, Huczyński A, Pyta K, Brzezinski B, Bartl F." Biological properties of Schiff bases and azo derivatives of phenols". Curr Org Chem 2009, 13: 124-148 .
- [14]. Xu,H. and Zeng,X. "Synthesis of diaryl-azo derivatives as potential antifungal agents ".Bio org.Med.Chem.Lett. ,(2010) ,20, 4193-4195 .
- [15]. Hatice K."Synthesis & spectral characterisation of novel azo-azomethine dyes ", Turk.J.Chem. , (1999) , 23,67-71 .
- [16]. Janeczek H, Sek D, Ostrowski S. " Similarities and differences between azomethines and ketimines " , Spectrochim Acta A Mol Biomol Spectrosc, (2007) , 66,1030-41.
- [17]. Meghasham N. N. "Synthesis, Characterization & Antimicrobial Activities of Substituted 2, 5- Disubstituted Phenyl-6-Substituted Phenyl Sulfonamide / Azomethine / Azo. ", IJARCS ,( 2014) , 1, 70-86 .
- [18]. S. Patai, Ed., "The Chemistry of the Carbon-Nitrogen Double Bond", J. Wiley & Sons, (1970), London .
- [19]. Muhammad A. A., Karamat M., "Synthesis, Characterization and Biological Activity of Schiff Bases" , IPCBEE ,(2011) , 10 , 1-7 .
- [20]. C. J. Patil et al. , " Synthesis & characterization of ketimines from 1-acetyl naphthalene with derivatives of aniline " , Der Pharma Chemica, (2016) , 8 (1) , 99-103 .
- [21]. C. J. Patil and C. A. Nehete , " Synthesis & characterization of ketimines from 5-Chloro-4-methyl-2-hydroxy-acetophenone with disubstituted anilines " , Der Pharma Chemica, (2015) , 6(10) , 1-6 .

- [22]. A. Saxena , "Synthesis and characterization of schiff base salicylaldehyde and thiohydrazones and its metal complexes " , Pelagia Research Library, (2013) , 4(4):152-154 .
- [23]. Prakash, C.R., Raja, S., "Synthesis and Antimicrobial Activities of Some Novel Schiff Bases of 5-Substituted Isatin Derivatives " , Rasayan Journal of Chemistry, (2009) , 2, 960- 968.
- [24]. SANDRA S. & DOO Aura. , " SYNTHESIS AND STRUCTURE OF VANILLIN AZOMETHINES" , CI&CEQ ,(2009) , 15 (4) 279–281 .
- [25]. R.K. Agarwal, S. Prasad, Bioinorg , "Synthesis, Characterization & Antimicrobial Activities of azo methines of 3-methoxy-4-butoxy benzaldehyde " , Chem. Appl. , (2005), 3, 271-288 .
- [26]. Furniss.S.Brian , Vogels textbook of practical organic chemistry " , 5th edition , John Wiley & Sons. Inc. , NY,(1989) .
- [27]. Abdullah J.Kadhim and Ali G.Kadhim" Synthesis & characterization some of  $\alpha$ ,  $\beta$  - unsaturated aliphatic ketones derivatives " , AL-Qadisiyah Journal for science, (2010) , 15 ,2, 45-62 .
- [28]. Bassar,M.A. & V.D.Phule , "Synthesis & antimicrobial activity of some new Schiff bases " , Orient.J.Chem. , (2000) , 16 , 553 .
- [29]. R.M.Silverstein , " Spectrometric identification of organic compounds" , 5th edition ,Wiley , NY ,(1991).
- [30]. Schraml, Jan; Bellama, Jon M. , " Two-Dimensional NMR Spectroscopy" New York: Wiley. , (1988) , pp. 28–33, 49–50
- [31]. Wadher.S.J. , Puranik M.P. &Karande N.A. , "Synthesis & Biological evaluation sciff base of dapsone &their derivative as antimicrobial agents", Inter.J.Pharm.Tech.Res. , (2009) , 1 , 22-33 .
- [32]. Jufe, G. S. "New hypnotics: perspectives from sleep physiology". Vertex (July–August 2007). 18 (74): 294–9 .
- [33]. Löscher, W.; Rogawski, M. A. "How theories evolved concerning the mechanism of action of barbiturates".(2012). Epilepsia. 53: 12–25.
- [34]. Huedo-Medina, T. B.; Kirsch, I.; Middlemass, J.; Klonizakis, M.; Siriwardena, A. N. "Effectiveness of non-benzodiazepine hypnotics in treatment of adult insomnia: meta-analysis of data submitted to the Food and Drug Administration"(17 December 2012) , BMJ. 345: e8343.